

This is a refereed journal and all articles are professionally screened and reviewed

ORIGINAL ARTICLE

## Necrotic Araneism. A Review of the *Loxosceles* Genus. I. General Aspects, Distribution and Venom Composition

<sup>1</sup>Hector Gabriel Ramos Rodríguez and <sup>2</sup>José D. Méndez

<sup>1</sup>School of Geography, Faculty of Philosophy and Letters, National Autonomous University of Mexico (UNAM).

<sup>2</sup>21<sup>st</sup> Century National Medical Center, Mexican Institute of Social Security, P.O. Box A-047, Mexico City, D.F., Mexico C.P. 06703.

Hector Gabriel Ramos Rodríguez and José D. Méndez,: Necrotic Araneism. A Review of the *Loxosceles* Genus. I. General Aspects, Distribution and Venom Composition, *Adv. Environ. Biol.*, 2(1): 9-19, 2008

### ABSTRACT

The “violin” spider, *Loxosceles sp* is implicated in causing necrotic skin lesions, secondary complications, and in some instances, even considered life-threatening. Fatalities are rare but are much more common in children, the ill, and the elderly. Diagnosis is often made by examining the lesion after the bite and rarely is a spider identified at the time of the attack. The “fiddle back” spider is distributed extensively in Mexico, with the majority of the species living in the Northern states. These spiders have a preference of hiding in closets, boxes, waste material, under rocks and in crevices and are considered “synanthropic”.

**Key words:** Arachnidism, *Loxosceles*, Necrotic arachnidism, Necrotic ulcer, “Violin” spider, Venomous spiders, Spider envenomation

### Introduction

The first representatives of the Arachnidea class appeared in the Silurian period of the Paleozoic era, over 360 million years ago. After acquiring aquatic habits, they reached their maximum development during the Carboniferous period of the same era, approximately 300 million years ago. It is during this period when they became terrestrial. The current species appeared during the Cenozoic era, 60 million years ago (Table I)[1,2].

Spiders, just as scorpions, are important to the public health sector since their bites are often harmful to adults[5]. They use their venom to kill insects and as a defense mechanism[6,7], but considering their dimensions, the amount of the inoculum and how fragile their oral organs are, it is

rare for them to cause severe lesions in subjects[8,9]. Nonetheless, of the more than 50,000 known species recognized worldwide, only 180 of them (0.45%) defend themselves aggressively and have chelicerae large enough to penetrate the human skin[10,11,14]. The species of greatest medical importance in the American continent are the “black widow” (*Latrodectus mactans*) and the “brown” (*Loxosceles sp*)[8,12]. In fact, the World Health Organization considers four genera of spiders to be of real medical interest due to the clinical manifestations they cause, and the lethality of their venom, three of them belonging to the infraorder Araneomorphae: *Latrodectus*, Theridiidae family; *Loxosceles*, Loxoscelidae family; *Phoneutria*, Ctenidae family. The fourth genus belongs to the Mygalomorphae infraorder (Atrax, Hexathelidae family)[13]. They are

### Corresponding Author

Dr. José D. Méndez, 21<sup>st</sup> Century National Medical Center, Mexican Institute of Social Security, P.O. Box A-047, Mexico City, D.F., Mexico C.P. 06703.  
Telephone Number: +52 55 5627 6900 ext. 21561  
Fax Number: +52 55 5227 6109  
E-mail: mendezf@servidor.unam.mx

**Table 1:** Taxonomy of the *Loxosceles* genera spider [3, 4].

|            |                              |
|------------|------------------------------|
| Kingdom    | Animalia                     |
| Phylum     | Arthropod                    |
| Subphylum  | Chelicerata                  |
| Class      | Arachnid                     |
| Order      | Araneae                      |
| Suborder   | Labidognatha                 |
| Infraorder | Araneomorphae (real spiders) |
| Family     | Sicariidae (Loxoscelidae)    |
| Genus      | <i>Loxosceles</i>            |
| Species    | <i>Loxosceles sp</i>         |

shy animals, never aggressive and only inflict bites when threatened, if violently touched or when pressed, almost always accidentally (70% during the night)[14,16]. One must remember, that spiders do not attack in groups, do not wait in lieu of becoming aggressive and some may even jump, but not intentionally attacking humans[17].

The incidence is unknown, but it is thought that females are more affected, and attacks usually occur in children[9,14,18]. Cases of intoxication by the venom of the *Loxosceles* in template zones of Europe, Africa, Australia, Asia and America have been recorded, each year registering an important number of cases of intoxication due to spider bites. In México, these observations were recorded in the states of Jalisco, Michoacán, Sinaloa, Sonora and Guanajuato as those of greatest incidence[19,20]. Some species are endemic, such as *L. tehuana*, unique to Oaxaca and Puebla[21]. In fact, close to 300,000 doses of sera are administered annually to save the lives of people that are stung or bitten by scorpions, snakes and black widow spiders [22,23]. Surely, the exposure to the *Loxosceles* venom can be mortal, be the cause of absenteeism from work for several weeks or leave important secuelae from an esthetic standpoint[24].

It is the second most frequent spider to cause accidents in countries such as Mexico and Peru[25].

### Epidemiology

Loxoscelism was first described in 1872 when Chilean physicians associated a peculiar skin lesion known as “a gangrene ulcer from Chile” or “gangrene in flame” to bites from *Loxosceles laeta*[26,27]. In the United States, *Loxosceles reclusa* was the first spider related to necrotic arachnidism (named by some as chronic arachnidism[28]) in 1957[29,30]. In 1984, at least five deaths were reported[31]. Nonetheless, severe cases started occurring at the end of 1800, and currently in endemic areas, this spider continues to be of clinical importance[32]. *L. laeta*, that produces more severe reactions than *L. deserta*, was imported from South America to Los Angeles previous to 1969[33]. Mortality is seen in children under seven years old in the Midwest and Southeast[34]. It is said that Caucasian individuals are the most sensitive to the venom[35].

The *Loxosceles* genus in America is represented by over 90 species, which are reported as agents causing loxoscelism to *L. laeta*, *L. spadicea*, *L. rufescens* and *L. gaucho* for South America, while for the United States and northern Mexico they are *L. reclusa*, *L. arizonica* and *L. deserta*. However, some authors consider that all species are capable of producing necrotic arachnidism or loxoscelism[36]. In 1997, the American Association of Poisoning Control Centers registered close to 2,000 bites by *L. reclusa* (endemic in Alabama), that occurred largely during the summer months when the spiders are more active[37].

In Argentina, accidents due to *Loxosceles* rank second due to spiders and represent 4% of the total of venous animals. The majority of the arachnoids are found in home environments (57%), while the remaining 43% are located around homes[38]. In Chile, bites by poisonous animals are recognized as work accidents. Spiders are often found in warehouses, libraries, laundry mats, cleaning companies, supermarkets, clothing and hardware stores, factories, constructions sites and places adequate for their presence (food and refuge). The number of men bitten is greater than that of women[39]. In Central America and South America, *L. laeta* is endemic, and known to be the most toxic, dangerous and often responsible for several deaths a year[32,40].

In Brazil, of the 153 cases studied, 118 (71%) had skin lesions and 35 (22.9%) had visceral-skin lesions[41]. The majority of the patients were bitten in the arm or legs (29.4% in the arm and 45.7% in the leg). Since 1990, the number of people bitten by *L. intermedia*[79] increased. At pediatric hospitals in Peru, during 1964-1980, the greatest incidence occurred in babies, pre-scholars and boys, with bites seen mainly in the chest area and extremities[42, 43].

At the Mexican Institute of Social Security (IMSS), the violinist spider is not included in the list of poisonous animals that have driven almost three thousand people to the Emergency Room in the last three years, even though it is considered a dangerous arthropod[58]. *L. boneti* is a widely distributed spider in Mexico and one of the main species involved in poisoning humans[44].

### Zoogeography

The *Loxosceles* genus includes at least 56 species in Eurasia, Africa and America; 54 in the American continent, one in the Mediterranean region and another in South Africa[29]. The most commonly known, *L. reclusa* and *L. laeta* are located in North, Central and South America. It is possible that the extensive geographic location of certain species occurred due to their distribution in commercial articles[45,46,47]. In fact, they are widely distributed in regions with tropical and warm climates. The

preferred climates identified are: temperate forest, rainy forest, grasslands (savanna), chaparral, and desert or dunes[31]. Mascaro (1974) mentions that *L. laeta* can be found extending from the United States down to the Patagonia[48]. *L. deserta* is mainly found in the Mojave and Sonora deserts, in the San Joaquin Valley and in areas adjacent to Mexico[49,17]. Cases of loxoscelism have been reported in Veracruz, Puebla, Morelos[7] and Mexico City[48]. Hoffman (1976) and Gerstch (1983) point to the entomogeography of the 38 species recognized in Mexico, of which 30 (78%) are endemic (Map 1)[50,51,52]. *L. deserta* has been reported in North and South Baja California[53]. However, until now it has not been possible to associate certain species with detected cases of necrotic arachnidism (arachnidism, araneism or arachnoidism)[54] since the majority of the times the spider remains unidentified. In Chiapas, *L. tehuana* is endemic. It is considered the most poisonous spider and the majority of the people bitten live in rural areas[55].

Similarly, *L. reclusa* is known to live on the Pacific coast of Mexico, extending from the Southern part of Sinaloa to Central America, inhabiting warm-humid and semi-dry zones[15,56]. The *L. devia*, *L. candela*, *L. belli* species have been identified. *L. devia* is the most widely seen, followed by *L. belli* and *L. candela*. *L. belli* registers in first place in the state of Nuevo León. There is a sympatric relationship between *L. devia* and *L. candela*, in addition to *L. belli* and *L. candela* for the municipalities of Monterrey and Garcia, respectively[36].

In Brazil, *L. gaucho*, *L. laeta* and *L. intermedia* are abundant with important levels of household infestation by the latter and mortality rates of 0.2%[16,57]. In the central zone, they are found in 41% of urban households and 24% of rural homes[58]. The age groups most affected are the first five decades. They have been found in hospital settings and academic areas[59].

In Spain, *L. rufescens* is classified as subtrogliphilic, that is to say, it lives part of the year refuged in caves and when outside, it is often found under rocks or in shady places[60].

### Morphology

The *Loxosceles* genus, to which *L. reclusa* belongs (“brown recluse spider”) and *L. laeta* (“South American brown spider”), are animals with bodies measuring 7-15 mm long and 0.2 to 2.5 cm in diameter dependent on diet and habitat (*L. laeta* measure 3-4.5 cm) [61, 30, 62]. *L. spiniceps* has a long body 8-10 mm and legs measuring 25-50 mm[63,64]. They possess an external skeleton composed primarily of chitin and a body divided into two regions or tagmas[13]. The prosome is light brown or gray and the opistosome is cream olive

colored (covered with fine hairs), linked by a structure called a pedicle (Figure 1)[65].

*L. arizonica* varies in color from yellow to dark brown[66]. Both genera reach a weight of 0.5 g. They have six simple eyes of equal size, a diade at the front of the cephalothorax and two others towards the back and to the side of the first, forming a triangle[56,59,67]. Their vision is excellent[68]; however they are the only spiders that cannot see from behind since they do not have the backend eyes (Figure 2)[69]. The legs measure 0.8-4 cm and are covered with fine hairs (with tactile function)[70] and have nails on the tarsus, longer than those of the males. The darker and more sclerosed part on the back of the cephalothorax includes a central anterior region in form of a violin, the “neck” pointing towards the abdomen[17] from where the name “violin spider” or “fiddle back spiders” derives[8,45,71] (Figure 1). In the *L. devia*, there is variation in intensity and in certain species; it is difficult to identify[49]. The majority is very hairy[72].

The males differ from the females in their sac-form expansion seen in the distal segment of the pedipalpi (stridulatory apparatus) or tactile organs on the sides of the chelicerae, that correspond to the seminal receptacle or spermatheca sac[58,67]. Each pedipalpe has six segments[13]. The Labidognatha suborder is characterized by its chelicerae (inoculating teeth) horizontally placed and when biting, cross over like a tweezers (Figure 3)[70].

### Biology and ethology

These spiders have night habits or are crepuscular (lucifugous), sedentary and segregary[75,76] allowing them to catch insects and other arachnids[14,45,56]. Its movements are rapid in comparison to other spiders[73]. In Argentina, the following results were obtained: 0% in the morning, 14% in the afternoon and 86% are seen during the evening[38]. They live conforming dense populations (up to 2,055 collect in a single house)[77], however it has been seen that *L. laeta* and *L. intermedia* do not share the same microhabitats[34,78]. They are often found under tree barks, in stored wood and bricks, under rocks, in livestock facilities, in bamboo plants[79], caves and museums. In houses (synanthropic environment[75]), where they mainly arrive in the fall, they are often found in dark, hidden places such as bathrooms, under furniture, over blinds, attics, boxes, clothes, bed sheets, walls (preferably corners), behind frames, mirrors, dents and abandoned waste. This cryptozoic characteristic allows them to install and proliferate abundantly[80]. The majority of the attacks occur while victims are sleeping (38%) and when they are dressing (32%)[14,18,42,45]. The intradomiciliary accident is more frequent at night throughout the year[72] and

for *L. intermedia*, mainly during the months of October to March and between January and April in Peru[81,82], and in the South American cone from December to May[22]. They hibernate all winter and the majority of bites occur between March and October[16,43,61]. The majority of the spiders rarely inflict more than a bite. Nonetheless, it is known that a woman was bitten three times by the same spider[83]. In general, females are more dangerous, particularly during mating season and during the period when they are raising their offspring (May-August)[15,84], related to greater weight and dimensions in comparison to males, differences not found between *L. spinulosa* and *L. parrami* of medical importance in South Africa[72,85]. In addition, as with some poisonous snakes, spiders sometimes inflict "dry" bites inoculating little or no venom, a factor that influences in the victim's response[86].

In general, mating takes place during June and July[30]. For *L. intermedia*, females have immature gonads during the months June, July and August[81]. They are more active at temperatures between 4° C and 30° C, although they may tolerate freezing temperatures of over 100 °F. *L. intermedia* withstand higher temperatures, while *L. laeta* tolerate lower ones[87]. They are oviparous producing up to 300 eggs during their reproductive life with an incubation period of 20-36 days[67,84,88]. They grow while shedding skin (exubia), a process known as ecdysis[67]. They reach maturity at a year, after 7-12 sheds[67,70] and a longevity of 2-4 years[7,49,56,89]; and more than 7 under experimental conditions and surviving 6-12 months without consuming water or food[30,67,90]. Life expectancy is greater in unmated animals[70].

They are basically insectivores: flies, moths, crickets, cockroaches, small arthropods and other bland body animals[17], although occasionally they feed on decomposing animals. The nymphs[91], youngsters and adults practice cannibalism[67,70,92]. *L. reclusa* has a predilection for dead animals[93]. In Brazil, *L. amazonica* shares a day refuge with small insects (ants and larvae), in places with little light and high humidity; *L. reclusa* prefers dry and damp places, while *L. rufescens* like fresh and humid hideouts[52,75,94]. Their web is irregular, lax and dirty similar to a bundle of cotton [89], without a specific pattern that serves as a refuge during the day[42,76].

The economic importance for mankind lies on the control of many destructive insects and plagues[31]. Health benefits are the consumption of hematophagous insects that are disease transmitters[95]. Biting insects such as flies, ticks, bedbugs and killer beetles, among others, may cause wounds similar to those produced by a bite by *L. reclusa*[33].

### Venom composition

Few spider venoms have been studied in detail, but it is known that they use it to immobilize being mainly cytotoxic and eat their catch[8,12]. The *Loxosceles* venom is composed of at least 10 or 12 proteins (similar to that of the rattle snake). It is thermo-labile and its enzymatic activity is superior to that of the *Latroectus*[9,58,96]. It is rich in enzymes: hydrolases, lipases, peptidases, collagenases, alkaline phosphatase, 5-ribonucleotidase, phosphohydrolases, proteases, hyaluronidase, acid phosphatase and phospholipase A<sub>2</sub>. The last three have great antigenic ability, and therefore great sensitivity[97,98]. The proteolytic effects are seen at the level of the constituents of the extracellular matrix such as fibronectin, fibrinogen, entactin, and heparin-sulphate proteoglycans[99]. Some authors consider it more potent than that of the rattle snake and cobra, since amounts of 0.5 µg (no more than 30 µg of protein) cause dermal necrosis in rabbits[34,40,100]. In fact, the bite causes a superficial intradermal inoculation of about 0.3 ml of venom, 0.5 to 1 µl of *L. laeta* and *L. rufipes* and 0.25-0.62 µl (0.36 µl on the average) for *L. reclusa*, in general, the male produces less but the amount that the inoculum is toxic[1,101,102]. It has necrotizing, hemolytic, vasculytic and coagulating[58]. It produces local edema, a thickening of the endothelium, accumulation of inflammatory cells, intravascular coagulation, thrombocytopenia and hemorrhaging and necrosis. The generalized systemic symptoms such as fever, malaise, pruritus and exanthema are common, while hemolysis, intravascular coagulation, thrombocytopenia and renal failure happen in approximately 16% of the victims[103]. Nonetheless, even large amounts of the venom from *L. gaucho*, *L. laeta* and *L. intermedia* have low proteolytic, myotoxic and phospholipase A<sub>2</sub> activity levels[104]. After being inoculated, the venom is more rapidly disseminated lymphatically than via the blood stream[56]. It is often light colored, of viscous consistency and contains enzymes with hemolytic, gelatinolytic, fibrinolytic and fibrinogenolytic activity in mammals (Table 2)[14,57,105]. The proteolytic action dependent on calcium on band 3 of the transmembrane protein of the erythrocytes makes them sensible to the activation of the complement with intravascular hemolysis following[106]. Hialuronidase (known as a "disseminating factor"), the nucleotide similar to adenosine triphosphate and citrate are the major components of the spider venom. The hialuronidase and the citrate (anticoagulant, chelating factor and buffer) are considered active components, and recently, extracellular ATP has been considered as toxic for certain cells[10,52]. Enriched with biogenic amines, not conceptualized within toxic fractions, they are

**Table 2:** Elements involved in the physiopathology caused by *Loxosceles* poisoning

| Venom contents   | Induced by the venom                            |
|--|---|
| Phosphodiesterases   | Histamine                                       |
| 5' ribonucleotide phosphohydralase                               | Serotonin                                       |
| Sphingomyelinase B and D   | Complement activation                           |
| Alkaline phosphatase   | Bradikinin                                      |
| Phospholipase A  | Prostaglandins                                  |
| Hyaluronidase  | Fibrinogen                                      |
| Collagenase  | Activation of V and X factors                   |
| Elastase   | Prothrombin                                     |
| L-amino acid – oxidases  | Thromboxan B <sub>2</sub>                       |
| Deoxyribonucleases   | Chemotactic substances                          |
| Adenosyntriphosphatase   | Interleukin 8                                   |
| Acetylcholinesterase   | Proteases                                       |
| Metalloproteinases   | Growth Factors                                  |
| Polyamines (spermine)  | Changes in the adhesion proteins                |
| Gamma aminobutyric acid  | Activation of polymorphonuclear cells           |
| Histamine  | Activation, expression and secretion of T cells |
| Serotonin  | Alpha oncogen related to growth                 |
| Neurotoxic peptides  | Monocytic chemotactic protein                   |
| Levarterenol bitartrate  | Degradation of erythrocytic glucophorins        |
| Antibactericide substances*                                      | * not proven                                    |
| Neurotoxin (glycosilated protein with sphingomyelinase activity) |   |




**Fig. 1:** Left: Female *Loxosceles* sp located in the state of Jalisco. Right: A) Female and B) Male of *Loxosceles* sp, from the state of Quintana Roo (Property of the National Collection of Acarus of the UNAM Institute of Biology, México).

capable of producing intense pain. Spermine may represent the final toxic product of the specialization that has occurred in certain lines of spiders during their evolution[105]. In fact, the polyaminamides, polar compounds of low molecular weight, act at the membrane stabilization level, blocking Ca<sup>2+</sup> and Mg<sup>2+</sup> channels, chelation of metallic ions and presynaptic and postsynaptic transmitters[107]. The acylpolyamines of low molecular weight are paralyzing and are composed of an aromatic portion containing amino acids and polyamines. These substances are main components of many poisonous spiders and are named “glutamate receptor antagonists”[10], rich in proteins with N-mannase residues (Loxolisine B) that provide the gelatinolytic activity, identified as two zymogens (serine-protease) of 85 and 95 KDa with no laminolytic activity[99,108]. The glycosilation of certain loxolisine B residues, a glycoprotein of 32-35 KDa, with metalloproteinase activity (endopeptidase containing an atom of zinc and secreted by connective tissue cells, phagocytes and a number of transformed cells) is important for the dermonecrotic activity. The substrates are collagen, fibronectin, elastin and alpha-1-antitripsin, among



**Fig. 2:** Left: Electron Microscopy Microphotograph, frontal view of the cephalothorax showing the ocular formula (2, 2, 2) and the stridulatory apparatus[73,74].

others[13,109,110]. The fibronectinolytic and fibrinogenolytic activities of the venom from *L. intermedia* are produced by a metalloproteinase of 20-28 KDa (Loxolisine A)[99,110]. The bite by



**Fig. 3:** Position of the inoculating hooks (arrows). A: Araneomorphae infraorder (*Phoneutria* spp), B: Mygalomorphae infraorder (*Tarantula*)[13].

*L. reclusa* causes intravascular necrosis associated to spherocytosis, changes that persist for several days to a week[14,111].

Sphingomyelinases B and D (or phospholipase D)[84,112] are the most important skin necrotic factors. They adhere to cell membranes causing platelet aggregation, erythrolysis, damage to blood vessels, fibrinogenolysis and an Arthus type reaction[8,71,108,113]. This is due to the events that involve sphingomyelinase metabolites including the inhibition of proliferation, differentiation and cell growth[40,112] since it induces the expression of metalloproteinase 9. Similarly, the venom induces apoptosis of human keratinocytes when increasing the expression of Metalloproteinases 2 and 9. In addition, the activation of these cells causes the production of TNF- $\alpha$ , a cytosine involved in local inflammatory processes. The venom from *L. deserta* induces the expression of the vascular endothelial growth factor in keratinocytes, which causes angiogenesis and vascular permeability contributing to vasodilatation, edema and erythema[35,114,115]. Platelet aggregation is a product of the adherence of the amyloid P serum glycoprotein to the platelet membrane in the presence of calcium ions, which triggers disseminated intravascular coagulation. Sphingomyelinase catalyzes the hydrolysis of sphingomyelin and induces the intrinsic activity of lisophospholipase D towards lisophosphatidylcholine producing lisophosphatidic acid, known to be an inducer of platelet aggregation, endothelial hyperpermeability and pro-inflammatory responses[40,112]. This enzyme has dermo-necrotic action in rabbits, and is toxic in mice and horses[52]. The secretion of serotonin by activated platelets induces chemotaxis of polymorphonuclear cells towards the lesion site, contributing to local vasculitis and necrosis[13,42]. The necrotizing activity is not associated to low molecular weight fractions such as inosine and nucleotides. Geren (1975) isolated two 34 KDa proteins that seem to be responsible for toxicity[105]

and the allergenic capacity[56]. Toxin I cause characteristic lesions and is lethal for rabbits and mice. Toxin II is not toxic for mice, is lethal for rabbits but does not cause necrotic lesions[105].

Currently three isoforms ( $P_1$ ,  $P_2$  and  $P_3$ ) are known with a relative mass of 28-35 KDa.  $P_1$  hydrolyzes sphingomyelin with lesser efficiency than  $P_2$ , that may be attributed to a substitution in the 203 position (Pro-Leu) and local substitutions of amino acids in the hydrophobic channel for the recognition and linking of the substrate. The third is inactive[76,116,117]. The sequence of 305 amino acids in the sphingomyelinase of *L. reclusa* is identical in 87, 85 and 60% to that of *L. arizonica*, *L. intermedia* and *L. laeta*, respectively; and 84 and 46% to SMasaD 1 and 2 of *L. boneti*[112]. In addition, crossed reactivity studies suggest that sphingomyelinase D (31-35 KDa) originated in the ancestral lineage of *Loxosceles/Sicarius*. It is inferred that the bites of all of the species from this group are capable of producing dermonecrotic lesions. The response between *L. boneti* (Mexico) and *L. laeta* (South America) is attributed to the filogenetic correlation due to its geographical origin[118]. These proteins vary in molecular mass among species: 31-32 KDa for *Loxosceles* from North America, 32-33.5 for the African type and 32-33 for *Sicarius*[119].

In *L. intermedia*, the hemolytic activity dependent on the complement and the necrosis inducing factor are associated to a 35 KDa protein, known as F35 with a potent lethal effect on mice. This is generated from the third larvae stage on and increases in quantity and toxic activity as the spider reaches the adult stage which coincides with the appearance of the abilities to disseminate, hunt and defend itself deemed necessary for the survival of these spiders[16,120]. A pronounced seasonal variation is seen in *L. laeta* in relation to the poisonous activity[121] considered more toxic than *L. reclusa*[4,15,52]. The venom of this species induces the production of alpha and beta kinases as the expressed and secreted regulator of the activation of normal T lymphocytes (RANTES), the monocyte chemo-attracting protein I (MCP-1), interleukin 8 (IL-8) and the alpha oncogene related to growth (GRO- $\alpha$ ) that probably participate in the appearance of dermonecrosis mediated by neutrophils and T lymphocytes. In addition, the activation of the complement causes the release of C5b, neutrophil chemo-attractants that generate tissue damage at the site of the bite[13,42]. Similarly, they cause the rupture of glycoporphins from the erythrocytic surface, easing the activation of the complement and hemolysis[121].

In Spain, *L. rufescens* is less poisonous than *L. laeta*[120]. However, it is not a remote possibility that other species cause these types of lesions [54]. In fact, the degree of systemic poisoning varies



Map. 1: Loxosceles species in Mexico (Hoffmann, 1976 and Gertsch, 1983)

between species[29]. The lethal dose (LD<sub>50</sub>) of *L. intermedia* is 0.48 mg/kg that of *L. gaucho* 0.74 mg/kg, while for *L. laeta* is 1.45 mg/kg. The dermonecrotic and lethal activities are detected in the A fraction of the three species, whose largest protein is 35 KDa in *L. gaucho* and *L. intermedia*, but 32 KDa (reported as sphingomyelinase) in *L. laeta* responsible for necrotic damage. All contain common residues in their N-terminal sequences. The identity is 50% between *L. gaucho* and *L. reclusa* and 61.1% between *L. intermedia* and *L. reclusa*[10,104]. Ontogenic differences are seen in the venom action in *L. intermedia*, being more lethal and dermonecrotic that of adult spiders than those from newborn, and of greater potency in females[13]. The volume that they inject when biting is in direct relation to the time that has passed since they last ate, with greater availability of venom if submitted to prolonged fasting[121].

On the average, they produce 0.1 mg of venom (0.15-0.45 µl) by electrical stimulation with 1.5 volts on the cephalothorax[13,42,76], place where the poisoning apparatus is located, composed of continuous apocrine glands, covered by secretory epithelium and surrounded by muscle cells. From *L. laeta* we can obtain 60 µg, while from *L. intermedia* and *L. gaucho* 30-40 µg[106]. From dissected glands, 0.8 µg of toxin can be obtained[120,121].

In conclusion, this paper provides general information related to the “violin” spider, *Loxosceles sp.*, its distribution in México and other countries, thus as the venom composition and its effects and cosequences in humans and animals.

References

1. Martino, O., T. Orduna and M. Espinosa, 2002. La agresión de los animales y su importancia en la salud. [www.aps.org.ar/\\_private/La agresion \(Parte I\).pdf](http://www.aps.org.ar/_private/La_agresion_(Parte_I).pdf).
2. Chemello E. Aranhas, acróbatas e perigosas. [www.ucs.br/ccet/defq/naeq/material\\_didactico/txtos\\_interativos\\_37.htm](http://www.ucs.br/ccet/defq/naeq/material_didactico/txtos_interativos_37.htm)
3. Platnick, N., 1993. Advances in Spider Taxonomy 1988-1991. With Synonymies and Transfers 1940-1980. USA: New York Entomological Society, American Museum of Natural History.
4. Cloudslay J.L., 1993. In: Medical Insects and Arachnides. Lane R P, Crosskey R W (Eds). Great Britain: Chapman and Hall.
5. Ramos, R.H. and R.I. Vázquez, 1999. Arañismo ocasionado por especies del género *Loxosceles* (Araneae; Sicariidae). *Rev Educ Salud* 29: 25-34.

6. Goodard, J., 1993. Physician's Guide to Arthropods of Medical Importance. USA: CRC Press.
7. Hunt, T.K. and E. Jawetz, 1989. Intoxicaciones en: Diagnóstico y Tratamiento Quirúrgicos. Way L W (Ed). México: Manual Moderno.
8. Moye de Alba, C.E., 1997. En: Manual de Terapéutica Médica y Procedimientos de Urgencia. Aguilar S C, Prado C E, Cañedo C S, Chevoile R J, Saucedo J, Kaplan S.M., Díaz A, Araña G J (Eds). México: McGraw Hill-Interamericana.
9. Russell, F.E., 1987. En: El Manual Merck. Berkow R, Fletcher A). España: Ediciones Doyma.
10. Odell G.V., H. Clement, L. Possani and A. Alagón, 2001. Spider Venom Toxins. J Venom Anim Toxins, 7: 335.
11. Biagi, F., 1974. Enfermedades Parasitarias. México: Manual Moderno.
12. Quintana, C.J.C. and P.R. Otero, 2002. Envenenamiento aracnídico en las Américas. MEDUNAB,5(13): 1-9.
13. Maguire, J.H. and A. Spielman, 1998. En: Principios de Medicina Interna. Vol. II. Fauci A S, Braunwald E, Isselbacher K J, Wilson J D, Martin J B, Kasper D L, Hauser S L, Longo D L. España: McGraw Hill-Interamericana.
14. Diniz, C.R., 1971. Venomous Animals and their Venoms. Vol. III. USA: Academic Press.
15. de Andrade, G.R.M., K.C. de Oliveira, A.L. Giusti, W.D. da Silva and D.V. Tambourgi, 1999. Ontogenetic development of *Loxosceles intermedia* spider venom. *Toxicon*, 37: 627-632.
16. Spiders Bite., 2002. Brown Recluse Spiders. Pest Management. University of California.
17. Berger, R.S., 1973. The unremarkable brown recluse spider bite. *JAMA* 225: 1109-1111.
18. Secretaría de Salud, 1999. Vigilancia Epidemiológica Semana 52, 1998. *Boletín Epidemiología*, 15: 18.
19. OPS-OMS., 1995. Clasificación Estadística Internacional de Enfermedades y Problemas Relacionados con la Salud (CIE - 10). Publicación Científica No. 554. USA: OPS.
20. Durán, B.C.G., 2006. Diversidad de arañas asociadas a viviendas de la ciudad de México. [www.estadis.eluniversal.com.mx/cultura/48817.html](http://www.estadis.eluniversal.com.mx/cultura/48817.html).
21. Editorial., 1998. Con Silanes, líneas de sueros contra venenos. *UNAM Hoy*, 33: 20.
22. Alagón, C.A., Anticuerpos terapéuticos: El caso de los antivenenos. Instituto de Biotecnología, UNAM. [www.ibt.unam.mx](http://www.ibt.unam.mx)
23. Araujo, M. and F. Otaiza, 2002. Efectividad del suero anti-*Loxosceles*. Síntesis de evidencia. ETESA.
24. Bioinformatics. Un veneno transgénico. UK, Oxford University Press, 2005.
25. Vest, D.K., 1999. The recluse spiders. Eagle Rock Research.
26. Luna, G.L., B.C. Ugarte, M.E. Soto and C.M.C. Ávila, 2000. Picadura por arácnido. *Rev Fac Med UNAM*, 43(3): 101-104.
27. Vest D.K., 1999. The Recluse Spiders. [www.srv.net/~dkv/recluse.html](http://www.srv.net/~dkv/recluse.html).
28. Weems, H.V., 2001. Brown recluse spider. University of Nebraska. Department of Entomology and Nematology.
29. Betten, J. and P. Garcia, *Loxosceles reclusa*. Brown recluse, the violin spider. Tehe University of Michigan - Museum of Zoology. Animal Diversity Web.
30. Arnold, T., 2002. *Loxosceles* in the United States. *eMedicine Journal*, 3(4): 2-11.
31. Spiders of Medical Importance., UCSD: Official web page of the University of California, San Diego; 2001-2002.
32. Keim, S.M., 2002. Spider bite, brown recluse. *eMedicine.com*.
33. Anjali, D., A.L. Heather and S.W. Jeffrey, 2000. *Loxosceles deserta* venom induces the expression of vascular endothelial growth factor (VEGF) in keratinocytes. *Inflammation*, 24(1): 1-9.
34. Solis-Rojas, C., J.J. Flores-Maldonado, W.A. Albeldaño-Vázquez, A. Flores Leal and K. Arèvalo Niño Distribución, 2002. del Género *Loxosceles* (Araneomorphae: Loxoscelidae) en el Estado de Nuevo León, México. Facultad de Ciencias Biológicas. Universidad Autónoma de Nuevo León, 2002.
35. Walley, S. and C. Johnston, 1997. *Loxosceles reclusa* y loxoscelismo: The Brown Recluse Spider and Envenomation Reactions. *Annals of Emergency Medicine*, 30: 28-32.
36. Borda, C.E., M.S. Delgado and M.J. Rea, 2005. Arácnidos de importancia médica de las especies *Loxosceles* en la región de influencia de la Universidad Nacional del Nordeste. CENPETROP (Resumen M-141).
37. Rodríguez, A.E. and R.M. Treviño, 2005. Riesgo laboral por mordedura de araña de los rincones. Departamento de Salud Ocupacional y C.A. Sección Riesgos Biológicos. Instituto de Salud Pública de Chile.
38. Murakami, M.T., P.M.F. Fernandes, D.V. Tambourgi and R.K. Arni, 2005. Structural Basis for Metal-Ion Coordination and the Catalytic Mechanism of Sphingomyelinas D. *ASBMB*, 1-26.
39. Mattosinho, S.G., U.M. Sezerino, M. Zannin, M. Grando, J.L.C. Cardoso, V.R.D. von Eickstedt and F.O.S. Franca, 1995. Geographic distribution of loxoscelism in Santa Catarina

- (Brazil) and species of *Loxosceles spp* involved and existents in the state. Departamento de Pediatria. Centro de Informaçõnes Toxicológicas.
40. Rodriguez, D.D., R.J. Mogollón, J. Guevara, A. Rodriguez, A.C. Eslava and M.F. Valdivia, Envenenamiento por arañas. Facultad de Medicina de la Universidad Privada "Antenor Orrego". Perú. [www.monografias.com/trabajos16/aracnidos-veneno/aracnidos-veneno.shtml](http://www.monografias.com/trabajos16/aracnidos-veneno/aracnidos-veneno.shtml)
  41. Araneismo, Aspectos epidemiológicos. Fundação Nacional de Saúde, 1999-2000.
  42. Estévez, J., J. Paniagua, R. Mancilla, A. Alagón, A. de Roodt, 2003. Experimental anti-*Loxosceles* ("brown spider") antivenom. [www.venenonemia.org](http://www.venenonemia.org).
  43. Harwood, R.F. and M.T. James, 1987. Entomología Médica y Veterinaria. México: Editorial Limusa.
  44. Condom, F., 19992. Manual de Terapéutica Quirúrgica. España: Editorial Salvat.
  45. Rook, 1989. Tratado de Dermatología. España: Editorial Doyma.
  46. Ramos, R.H. and R.L.M. Canul, 2000. Ulcera necrótica ocasionada por arañismo: un caso. Rev Educ Salud, 36: 7-10.
  47. Velter, R., 2000. Brown Recluse and Other Recluse Spiders. USA: IPM Education and Publication.
  48. Jiménez, J.M.L., 1996. En: Taxonomía y Biogeografía de Artrópodos de México. Hacia una síntesis de su conocimiento. Llorente B J, García-Aldrete A N, González-Soriano E (Eds). México: Instituto de Biología, CONABIO, Facultad de Ciencias UNAM.
  49. Gertsch, W.J. and F. Ennik, 1983. The Spider Genus *Loxosceles* in North America, Central America, and the West Indies (*Araneae, Loxoscelidae*). USA: Bull Am Museum Nat His.
  50. Cruz, W.L., Las arañas *Loxosceles*. <http://research.amnh.org/entomology/spiders/catalog/INTRO1.html>
  51. Berrian, J., 1997. Spiders of Baja California. San Diego Natural History Museum. [www.sdnhm.org/research/entomology/bcspider.html](http://www.sdnhm.org/research/entomology/bcspider.html)
  52. Hoffmann, A., 1976. Relación bibliográfica preliminar de las arañas de México (*Arachnida: Araneae*). México: Instituto de Biología, UNAM. Publicaciones Especiales.
  53. Vázquez, O.V., 1992. En: Arañas de Chiapas. Alvarez del Toro M (Ed). México: Universidad Autónoma de Chiapas.
  54. CUCUBA., 1999. Arañas Venenosas. México: Centro Universitario de Ciencias Biológicas y Agropecuarias. Universidad de Guadalajara.
  55. Braz, A., J. Minozzo, J.C. Abreu and I.C. Gubert, 1999. Chávez-Olórtegui C. Development and evolution of the neutralizing capacity of horse antivenom against the Brazilian spider *Loxosceles intermedia*. Toxicon, 37: 1323-1328.
  56. Barranco, C.O., Animales Ponzoñosos. La amenaza que viene. [www.tiempo.com.mx/not\\_detalle.php?id\\_n=5818](http://www.tiempo.com.mx/not_detalle.php?id_n=5818)
  57. Schenone, F.H., 2003. Cuadros tóxicos producidos por mordeduras de araña en Chile: Latrodectismo y loxoscelismo. Rev Méd Chile, 131: 437-444.
  58. Jiménez, C.A., Avance muy preliminar de la fauna de las cuevas de la Sebbética cordobesa. Grupo Espeleológico G40. [www.g40espeleo.com](http://www.g40espeleo.com)
  59. Wilson, C.D. and King E.L., 1999. Arthropod Bites and Stings. In: Dermatology in General Medicine. Vol. II. Freedberg I M, Eisen A Z, Wolff K, Austen K F, Goldsmith L A, Katz S I, Fitzpatrick T B (Eds). USA: McGraw Hill.
  60. Eliastam, M., G.L. Sternbach, B.M. Jay Manual de Medicina de Urgencias, 1994. Quinta Edición. España: Mosby/Doyma.
  61. Spiders are fun, but be wary of some. WildNet Africa (Pty.) Ltd. [www.wildnetafrica.com](http://www.wildnetafrica.com)
  62. Nieuwenhuys, E., 2001. Spiders an Immunology. University of Florida. Department of Entomology and Nematology.
  63. Hurtado, V.J.G., C.N. Sotelo and S.R. Ibarra, 2005. Envenenamiento por *Loxosceles reclusa* (araña "parda"). Rev Mex Pediatr, 72(2): 85-88.
  64. Bites and Stings. [www.ci.phoenix.az.us/FIRE/bitessna.html](http://www.ci.phoenix.az.us/FIRE/bitessna.html) 2002.
  65. Núñez, B.P., *Loxosceles sp.* OCTOLAB. Aracnotoxinas. [www.redtox.org:8080/Redtox/Templates/interiores\\_vis.dwt.jsp?idsubcat=ME](http://www.redtox.org:8080/Redtox/Templates/interiores_vis.dwt.jsp?idsubcat=ME)
  66. Brown Recluse Spider. [www.thebigzoo.com/Animals/Brown\\_Recluse\\_Spider.asp](http://www.thebigzoo.com/Animals/Brown_Recluse_Spider.asp)
  67. Insectos. [www.mayaparadise.com/rioenvs.htm](http://www.mayaparadise.com/rioenvs.htm)
  68. Parra, D., M. Torres, J. Morrillas and P. Espinoza, 2002. *Loxosceles laeta*, identificación y una mirada bajo microscopía de barrido. Parasitol Latinoam, 57: 75-78.
  69. Sánchez, J.A., 19993. Cien Años de la Zoología Médica. México: Instituto de Seguridad y Servicios Sociales de los Trabajadores del Estado.
  70. Echenique, G., 2003. *Latrodectus mactans* y *Loxosceles laeta*. Ponencia. Animales Venenosos.
  71. La araña de rincón, 2005. un peligro latente. [www.lacoctelera.com/premium-service](http://www.lacoctelera.com/premium-service).
  72. Suárez, H.M., A.J.J. Pérez, M.T. Rodríguez and M.E. Pena, 1998. Aracnoidismo sistémico. Rev Cubana Med Gen Integr, 14(1): 1-5.
  73. Neves A.H., B.A. Ribeiro, A.I. Sales, M.I. Rocha, 2004. Registro de *Loxosceles amazonica* Gerstch, 1967 (Araneae, Sicariidae) no Cariri Paraibano. Revista de Biología e Ciências da

- Terra 5(1): 1-4.
74. de Roodt A.R., O.D. Salomón, S.C. Lloveras and T.A. Orduna, 2002. Envenenamiento por arañas del género *Loxosceles*. *Medicina*, 62(1): 83-94.
  75. Vetter, R.S. and D.K. Barger, 2002. An infestation of 2,055 brown recluse spiders (Araneae; Sicariidae) and no envenomations in a Kansas home: Implications for bite diagnoses in nonendemic areas. *J Med Entomol*, 39(6): 948-951.
  76. Luciano, F.M. and N.J. Vasconcellos, 2005. Microhabitats occupied by *L. laeta* and *L. intermedia* (Araneae: Sicariidae) in Curitiba, Paraná, Brazil. *J Med Entomol*, 756-765.
  77. O mundo da aranha., 1974. [www.geocities.com/~esabio/aranha/aranha\\_principal.htm](http://www.geocities.com/~esabio/aranha/aranha_principal.htm).
  78. Goncalves, R.M. and T.D. Vilarinho, 2003. First record on *Loxosceles laeta* (Nicolet, 1849) (Araneae, Sicariidae) in the west zone of São Paulo City, São Paulo, Brazil, and considerations regarding its geographic distribution. *Rev Soc Bras Med Trop*, 36(3): 1-5.
  79. Ramires, E. P.F. Ferraz and S. Romao, 2002. Estudio preliminar do ciclo reprodutivo de *Loxosceles intermediata*. Universidade Tuiuti do Paraná.
  80. Maquiña, V.C., J.C. Hinojosa and R. Gutiérrez, 2004. Henríquez C, Ugarte C. Enfermedades por artrópodos. Parte I: Loxoscelismo cutáneo y cutáneo-visceral en el Perú. *Dermatol Perú*, 14(2): 1-10.
  81. Berry, H., 2001. Has the brown recluse spider reached Ontario?. *GardenWeb Forums*. [GardenWeb](http://GardenWeb.com).
  82. The spider – *Loxosceles reclusa*. Brown Recluse Spider Bite Site, 2002.
  83. de Oliveira, K.C., G.R.M. de Andrade, A.L. Giusti, W.D. da Silva and D.V. Tambourgi, 1999. Sex-linked variation of *Loxosceles intermedia* spider venoms. *Toxicon*, 37: 217-221.
  84. Spider bites. [www.dermatology.cdlib.org/cgi-bin/search.pl?query=Loxosceles%20envenomation&index=doj.swish](http://www.dermatology.cdlib.org/cgi-bin/search.pl?query=Loxosceles%20envenomation&index=doj.swish)
  85. Marchioro, C.A., M.L. Fischer and S.E. Marques, 2005. Desenvolvimento pós-embriônico de *Loxosceles intermedia* Mello-Leitão, 1934, *L. laeta* (Nicolet, 1849) e *L. gaucho* Gerstch, 1967 (Araneae; Sicariidae) criadas sob condições de alimentação monoespecífica. *Biotemas*, 18(1): 93-112.
  86. Lyon, W.F., 1991. Brown recluse spider. Ohio State University Extension Fact Sheet.
  87. Bücherl, W., 1969. Biology and venoms of the most important South American spider of genera *Phoneutria*, *Loxosceles*, *Lycosa* and *Latrodectus*. *Am Zool*, 157-159.
  88. Necrotic Arachnidism-Pacific Northwest, 1988-1996. *MMWR* 1996, 45(21): 433-436.
  89. Araneismo. [www.e-mas.co.cl/categorías/biología/sarna.htm](http://www.e-mas.co.cl/categorías/biología/sarna.htm)
  90. Spider Elimination. Brown recluse spider. [www.pestproducts.net/brownreclusebite.htm](http://www.pestproducts.net/brownreclusebite.htm) 2002.
  91. Sandidge, J., El veneno contra insectos puede empeorar las plagas de arañas. [www.axxon.com.ar/mus/info/030102.htm](http://www.axxon.com.ar/mus/info/030102.htm)
  92. Jacobs, S.B., 2003. Arañas comúnmente encontradas en Pensilvania (y dos especies raramente encontradas, pero con importancia médica). *Notas Epidemiológicas*. [www.ento.psu.edu/extension/factsheets/spanish/spidersSp.htm](http://www.ento.psu.edu/extension/factsheets/spanish/spidersSp.htm)
  93. Piola, J.C., Accidentes por arañas. Servicio de Toxicología del Santorio de Niños, Rosario, Argentina (2204).
  94. Swing, F.P., 1999. Acupuncture for treatment of a nonhealing brown recluse spider bite: a case report. *Medical Acupuncture a Journal for Physicians for Physicians*, 11(1): 1-6.
  95. Malaque, C.M., J.E. Castro-Valencia, J.L.C. Cardoso and F.F. de Siquiera, 2002. Clinical and epidemiological features of definitive and presumed loxoscelism in Sao Paulo, Brazil. *Revista do Instituto de Medicina Tropical de Sao Paulo*.
  96. Pastrana, J., R. Blasco and M.A. Pinillos, 2003. Picaduras y Mordeduras de animales. *Anales Hosp. Navarra*, 26(Supl. 1): 1-24.
  97. Villaveces, J.W., 1968. Gangrenous Spider Bite in Los Angeles County (Apparently by *Loxosceles reclusa*). *California Medicine*, 108(4): 305-308.
  98. Di Nardo, L.C., C.A. González and R. Bertrand, 2005. Picadura por arácnidos del género *Loxosceles*. *Boletín Epidemiológico*, 5: 6-8.
  99. Sanabria, H., A. Zavaleta, 1997. Panorama epidemiológico del loxoscelismo en el Perú. *Rev Peru Med Exp Salud Pública*, 14(2): 1-11.
  100. Tambourgi, D.V., F.C. Magnoli, V.L. Petricevich, M. Sucupira, Z.C. Benedetti, S.L. Assaf, E.W. Alveds, V.R.D. von Eickstedt, S. Jancar, W. Dias da Silva, 2002. *Fundação Butantan (Brasil)*.
  101. Barbaro, K.C., 1996. Biological, immunochemical and biochemical characterization of venoms of spiders of the genus *Loxosceles*. *Laboratório de Imunopatologia*. Instituto Butantan, Brasil.
  102. Blum, M.S., 1981. *Chemical Defenses of Arthropods*. USA: Academic Press.
  103. de O.B.O.C., M. Satake, K. Nonoyama and J.L.C. Cardoso, 2003. The calcium-dependent protease of *Loxosceles gaucho* venos acts preferentially upon red cell band 3

- transmembrane protein. *Braz J Med Biol Res*, 36(3): 309-313.
104. Blagbrough, S.I., E. Moya and S. Taylor, 1994. Polyamines and polyamine amides from wasps and spiders. *Biochem Soc Transactions*, 22: 888-893.
105. Veiga, S.S., W. Gremski, V.L.P. dos Santos, L. Feitosa, O.C. Mangili, H.B. Nader, C.P. Dietrich and R.R. Brentani, 1999. Oligosaccharide residues of *Loxosceles intermedia* (brown spider) venom proteins: dependence on glycosylation for dermonecrotic activity. *Toxicon*, 37: 587-607.
106. Aponte, D.M., C.O. Gutiérrez and J.M. Lozano, 2000. Detección de metaloproteinasas en cáncer gástrico y tejido normal mediante cimoografía. *Rev Colomb Gastroenterol*, 15: 170-175.
107. Monteiro, C.L.B., R. Rubel, L.L. Cogo, O.C. Mangili, W. Gremski and S.S. Vega, 2002. Isolation and identification of *Clostridium perfringens* in the venom and fangs of *Loxosceles intermedia* (brown spider): enhancement of the dermonecrotic lesion in loxoscelismo. *Toxicon*, 40: 409-418.
108. Rosse, W. and B.H. Franklin, 1998. En: *Principios de Medicina Interna*. Vol. I. Fauci A S, Braunwald E, Isselbacher K J, Wilson J D, Martin J B, Kasper D L, Hauser S L, Longo D L (Eds). España: McGraw Hill-Interamericana.
109. Lee, S. and K.R. Lynch, 2005. Brown recluse spider (*Loxosceles reclusa*) venom phospholipase D (PLD) generates lysophosphatidic acid (LPA). *Biochem J*, 391: 317-323.
110. Ruiz Maldonado, 1991. *Tratado de Dermatología Pediátrica*. México: Editorial Interamericana.
111. Paixão, C.D., C.W. van den Berg, F.M. de Freitas, R.M. Gonçalves and D.V. Tambourgi, 2006. Role of Matriz Metalloproteinases in HaCaT Keratinocytes Apoptosis Induced by *Loxosceles Venos* Sphingomyelinase D. *J Invest Dermatol*, 126: 61-68.
112. Malaque, C.M.S., M. Ori, S.A. Santos and D.R. Andrade, 1999. Production of TNF-alpha by primary cultures of human keratinocytes challenged with *Loxosceles gaucho* venom. *Revista do Instituto de Medicina Tropical de Sao Paulo*.
113. Hernández, J.C., R.J. Estévez and S.J. Paniagua, 2006. Araña violinista: problema médico en México. *Gaceta Biomédicas*. [www.biomedicas.unam.mx/presenta\\_gaceta.asp?fecha=10/06/2006](http://www.biomedicas.unam.mx/presenta_gaceta.asp?fecha=10/06/2006)
114. De Andrade, S.A., M.T. Murakami, D.P. Cavalcante, R.K. Arni and D.V. Tambourgi, 2006. Kinetic and mechanistic characterization of the sphingomyelinase D from *Loxosceles intermedia* spider venom. *Toxicon*, 47: 380-386.
115. Compositional variation and antigenic cross-reactivity among venoms of *Loxosceles* and their relatives. [www.venenonemia.org](http://www.venenonemia.org)
116. Binford, G.J., 2003. The phylogenetic distribution of sphingomyelinase D activity in venoms of haplogyne spiders. *Comp Biochem Physiol B*, 135(1): 25-33.
117. De Oliveira, K.C., G.R. de Andrade, A.L. Giusti, W.D. da Silva and D. Tambourgi, 1998. Sex-linked variation of *Loxosceles intermedia* spider venoms. *Toxicon*, 37(1): 217-221.
118. Schenone, H., T. Letonja and F. Knierim, 1975. Some data on the venom apparatus of *Loxosceles laeta* and the toxicity of its various animal species. *Bol Chil Parasitol*, 30: 37-4.
119. Tambourgi, D.V., B.P. Morgan, R.M. de Andrade, F.C. Magnoli and C.W. van den Berg, 2000. *Loxosceles intermedia* spider envenomation induces activation of an endogenous metalloproteinase, resulting in cleavage of glycoporphins from the erythrocyte surface and facilitating complement-mediated lysis. *Blood*, 95(2): 683-691.
120. Dueñas, L.A. and E.J.C. Martín, 1997. En: *Medicina Interna*. Tomo II. Texidor J R, Guardia M J (Eds.). España: Masson, SA.
121. Loxoscelismo. Araña del rincón (*Loxosceles laeta*). [www.2udec.cl/~lpalma/loxoscelismo/loxoscelismo.html](http://www.2udec.cl/~lpalma/loxoscelismo/loxoscelismo.html)